



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the same sort occur. Except for a few such, however, the botanist who knows the plants of an extended region will approve of all such unions as appear in a necessarily hasty review of the book.

In some few instances, though the editors have evidently done their work anew for this edition, and have not contented themselves with compiling from earlier editions or other sources, defective descriptions or the omission of really crucial characters are noticeable. Thus, the imbricate petals of *Anonaceæ* are still called valvate; *Potentilla rivalis*, var. *pentandra* is redescribed as having five stamens, whereas the number is usually six or eight, five being very exceptional in the specimens of the Engelmann herbarium and in many that have been examined in the field by Mr. Hitchcock; the petaloid filaments of *Thalictrum clavatum* are called club shaped, etc. Very useful distinctions between *Oxalis corniculata* and its variety *stricta* are afforded by the rhizomes and dichotomous inflorescence of the latter, from which *O. recurva*, which resembles it in some respects, differs in the trimorphic heterogony of its flowers. It might also have been well to note that the blue flowered flaxes, introduced in the East, belong to two well-marked forms, one of them, which has been separated under the name of *L. humile*, having widely-dehiscent capsules with ciliate septa, the other, with nearly closed capsules the septa of which are not ciliate. The reviewer must also plead guilty to having omitted the very important characters derivable in *Epilobium* from the innovations, which consist of sessile buds in no. 1, of dense rosettes at base of the stem in nos. 4 and 5, of running leafy shoots in nos. 7 and 8, of scaly rhizomes in no. 9, and of filiform bulbiferous shoots in nos. 2, 3 and 10. But whatever little defects may be noticed in one way or another, both amateurs and working botanists, who are concerned with the flora of the northern states anywhere east of the Rocky Mountains, will be grateful for so good a book, and feel disposed to congratulate the editors on the very satisfactory way in which they have brought it out.—
WILLIAM TRELEASE.

OPEN LETTERS.

Deep-water Nostoc.

In the sentence "When Dr. Wolle's 'Fresh Water Algæ' appeared, this Nostoc was not mentioned," in my note in the GAZETTE, November, 1889, p. 291, I referred to the deep-water Nostoc of Lake Michigan. If I had said "Our deep-water Nostoc was not mentioned," I should have expressed my thought more perfectly.

C. B. ATWELL.

Evanston, Ill.

Some more queer botany.

In a letter in this department last year a writer called attention to "some queer botany" which he found in a "doctor-book." If only it could be confined to this class of publications less harm would result than now